

## **Author Index**

Abel, T.W. and Rance, N.E.

Proopiomelanocortin gene expression is decreased in the infundibular nucleus of postmenopausal women (69) 202

Akira, S., see Yukawa, K. (69) 124 Alberch, J., see Canals, J.M. (69) 242

Alvarez-Maya, I., see Martinez-Fong, D. (69)

Arias-Montaño, J.-A., see Martinez-Fong, D. (69) 249

Arimatsu, Y., see Miyasaka, N. (69) 62 Arvidsson, U., see Ko, J.L. (69) 171

Baecker, P.A., Lee, W.H., Verity, A.N., Eglen, R.M. and Johnson, R.M.

Characterization of a promoter for the human glial cell line-derived neurotrophic factor gene (69) 209

Bannai, S., see Nakaso, K. (69) 155

Berezovska, O., Frosch, M., McLean, P.,

Knowles, R., Koo, E., Kang, D., Shen, J., Lu, F.M., Lux, S.E., Tonegawa, S. and Hyman, B.T.

The Alzheimer-related gene presenilin 1 facilitates notch 1 in primary mammalian neurons (69) 273

Bien, A., see Kreutz, M.R. (69) 232 Billings, L.M., see Schuller, J.J. (69) 149 Bing, G., see Feng, Z. (69) 144 Böckers, T.M., see Kreutz, M.R. (69) 232

Brown, L.A., see Poulter, M.O. (69) 44 Bustos, G., see Vecchiola, A. (69) 1

Canals, J.M., Checa, N., Marco, S., Michels, A., Pérez-Navarro, E. and Alberch, J. The neurotrophin receptors *trkA*, *trkB* and *trkC* are differentially regulated after excitotoxic lesion in rat striatum (69) 242

Cao, S., see Hu, X. (69) 35

Chang, R.C.C., see Feng, Z. (69) 144 Checa, N., see Canals, J.M. (69) 242

Ciani, E., Hoffmann, A., Schmidt, P., Journot, L. and Spengler, D.

Induction of the PAC1-R (PACAP-type I receptor) gene by p53 and Zac (69) 290 Collyer, P., see Vecchiola, A. (69) 1

Dragunow, M., see Woodgate, A. (69) 84 D'Souza, U.M., see Lammers, C.-H. (69) 281

Eglen, R.M., see Baecker, P.A. (69) 209 Elde, R., see Ko, J.L. (69) 171 Enikolopov, G., see Nishizaki, T. (69) 263 Feng, Z., Chang, R.C.C., Bing, G., Hudson, P.,
Tiao, N., Jin, L. and Hong, J.-S.
Long-term increase of Sp-1 transcription
factors in the hippocampus after kainic acid
treatment (69) 144

Figueroa, R., see Vecchiola, A. (69) 1 Frosch, M., see Berezovska, O. (69) 273

Goebel, D.J. and Poosch, M.S.

NMDA receptor subunit gene expression in the rat brain: a quantitative analysis of endogenous mRNA levels of NR1<sub>Com</sub>, NR2A, NR2B, NR2C, NR2D and NR3A (69) 164

Gong, C., see Yang, G.-Y. (69) 135 Gu, G., see Leng, N. (69) 73

Harper, P.S., see Wilkinson, F.L. (69) 10 Hatanaka, Y., see Miyasaka, N. (69) 62 Hoffmann, A., see Ciani, E. (69) 290

Hökfelt, T., see Johnson, H. (69) 21 Hong, J.-S., see Feng, Z. (69) 144

Honma, Y., Kanazawa, K.-y., Mori, T., Tanno, Y., Tojo, M., Kiyosawa, H., Takeda, J., Nikaido, T., Tsukamoto, T., Yokoya, S. and Wanaka, A.

Identification of a novel gene, OASIS, which encodes for a putative CREB/ATF family transcription factor in the long-term cultured astrocytes and gliotic tissue (69) 93

Hu, X., Cao, S., Loh, H.H. and Wei, L.-N.
Promoter activity of mouse κ opioid receptor gene in transgenic mouse (69) 35

Hudson, P., see Feng, Z. (69) 144 Hyman, B.T., see Berezovska, O. (69) 273

Inoue, M., Yamada, T. and Ueda, H.

Low dose of kyotorphin (tyrosine–arginine) induces nociceptive responses through a substance P release from nociceptor endings (69) 302

Ishii, T., see Nakaso, K. (69) 155

rons of aged rats (69) 21

Jarrott, B., see Sweerts, B.W. (69) 113
Jin, L., see Feng, Z. (69) 144
Jin, M.-h., see Miyasaka, N. (69) 62
Johnson, H., Hökfelt, T. and Ulfhake, B.
Expression of p75<sup>NTR</sup>, trkB and trkC in nonmanipulated and axotomized motoneu-

Johnson, R.M., see Baecker, P.A. (69) 209 Jones, A.L., see Wilkinson, F.L. (69) 10 Journot, L., see Ciani, E. (69) 290 Julien, J.-P., see Levayasseur, F. (69) 104 Kanazawa, K.-y., see Honma, Y. (69) 93
Kang, D., see Berezovska, O. (69) 273
Kawakami, H., see Miyachi, T. (69) 223
Kelley, M.R., see Zhou, F.C. (69) 186
Kiessling, M., see Sommer, C. (69) 53
Kimura, K., see Nakaso, K. (69) 155
Kitamura, T., see Miyachi, T. (69) 223
Kitayama, M., see Nakaso, K. (69) 155
Kiyosawa, H., see Honma, Y. (69) 93
Knowles, R., see Berezovska, O. (69) 273
Ko, J.L., Arvidsson, U., Williams, F.G., Law, P.Y., Elde, R. and Loh, H.H.
Visualization of time-dependent redistribution of δ-opioid receptors in neuronal cells

tion of δ-opioid receptors in neuronal cells during prolonged agonist exposure (69) 171 Koo, E., see Berezovska, O. (69) 273 Kreutz, M.R., Bien, A., Vorwerk, C.K., Böck-

ers, T.M., Seidenbecher, C.I., Tischmeyer, W. and Sabel, B.A.
Co-expression of c-Jun and ATF-2 charac-

terizes the surviving retinal ganglion cells which maintain axonal connections after partial optic nerve injury (69) 232

Labarca, R., see Vecchiola, A. (69) 1
Lammers, C.-H., D'Souza, U.M., Qin, Z.-H.,
Lee, S.-H., Yajima, S. and Mouradian, M.M.
Regulation of striatal dopamine receptors by
corticosterone: an in vivo and in vitro study
(69) 281

Law, P.Y., see Ko, J.L. (69) 171 Lawrence, A.J., see Sweerts, B.W. (69) 113 Lee, S.-H., see Lammers, C.-H. (69) 281 Lee, W.H., see Baecker, P.A. (69) 209

Leng, N., Gu, G., Simerly, R.B. and Spindel,E.R.Molecular cloning and characterization of

two putative G protein-coupled receptors which are highly expressed in the central nervous system (69) 73

Levavasseur, F., Zhu, Q. and Julien, J.-P. No requirement of α-internexin for nervous system development and for radial growth of axons (69) 104

Liu, X.-H., see Yang, G.-Y. (69) 135 Loh, H.H., see Hu, X. (69) 35 Loh, H.H., see Ko, J.L. (69) 171 Lorris Betz, A., see Yang, G.-Y. (69) 135 Lu, F.M., see Berezovska, O. (69) 273 Luna, J., see Martinez-Fong, D. (69) 249 Lux, S.E., see Berezovska, O. (69) 273

MacGibbon, G., see Woodgate, A. (69) 84 Magendzo, K., see Vecchiola, A. (69) 1 Manilal, S.B., see Wilkinson, F.L. (69) 10
Marco, S., see Canals, J.M. (69) 242
Marshall, J.F., see Schuller, J.J. (69) 149
Martinez-Fong, D., Navarro-Quiroga, I., Ochoa, I., Alvarez-Maya, I., Meraz, M.A., Luna, J. and Arias-Montaño, J.-A.
Neurotensin-SPDP-poly-L-lysine conjugate: a nonviral vector for targeted gene delivery to neural cells (69) 249

a nonviral vector for targeted gene delivery to neural cells (69) 249

Maruyama, H., see Miyachi, T. (69) 223

Matsuoka, T., see Nishizaki, T. (69) 263

Matsuoka, T., see Nishizaki, T. (69) 295

McLean, P., see Berezovska, O. (69) 273

Meraz, M.A., see Martinez-Fong, D. (69) 249

Meuser, T., see Pietruck, C. (69) 286

Michels, A., see Canals, J.M. (69) 242

Miyachi, T., Maruyama, H., Kitamura, T.,

Nakamura, S. and Kawakami, H.

Structure and regulation of the human NeuroD (BETA2/BHF1) gene (69) 223

Miyasaka, N., Hatanaka, Y., Jin, M.-h. and Arimatsu, Y.
Genomic organization and regulatory elements of the rat latexin gene, which is expressed in a cell type-specific manner in both central and peripheral nervous systems (69) 62

Mori, T., see Honma, Y. (69) 93 Morris, G.E., see Wilkinson, F.L. (69) 10 Mouradian, M.M., see Lammers, C.-H. (69) 281

Nakamura, S., see Miyachi, T. (69) 223
Nakashima, K., see Nakaso, K. (69) 155
Nakaso, K., Kitayama, M., Ishii, T., Bannai, S.,
Yanagawa, T., Kimura, K., Nakashima, K.,
Ohama, E. and Yamada, K.
Effects of kainate-mediated excitotoxicity
on the expression of rat counterparts of
A170 and MSP23 stress proteins in the
brain (69) 155

Navarro-Quiroga, I., see Martinez-Fong, D. (69)

Neal, J.W., see Wilkinson, F.L. (69) 10 Nikaido, T., see Honma, Y. (69) 93

Nishizaki, T., Matsuoka, T. and Sumikawa, K. Positive- and negative-regulation of Ca<sup>2+</sup> entry through ACh receptor channels by phosphorylation (69) 295

Nishizaki, T., Nomura, T., Matsuoka, T., Enikolopov, G. and Sumikawa, K. Arachidonic acid induces a long-lasting facilitation of hippocampal synaptic transmission by modulating PKC activity and nicotinic ACh receptors (69) 263

Nomura, T., see Nishizaki, T. (69) 263

Ochoa, I., see Martinez-Fong, D. (69) 249 Oertel, W.H., see Sommer, C. (69) 53 Ohama, E., see Nakaso, K. (69) 155 Patel, T.D., see Zhou, F.C. (69) 186
Pérez-Navarro, E., see Canals, J.M. (69) 242
Pierce Palmer, P., see Pietruck, C. (69) 286
Pietruck, C., Xie, G.-X., Sharma, M., Meuser,
T. and Pierce Palmer, P.
Multiple splice patterns of cyclic AMP response element-binding protein mRNA in the central nervous system of the rat (69)

Poosch, M.S., see Goebel, D.J. (69) 164
Poulter, M.O. and Brown, L.A.

Transient expression of GABA<sub>A</sub> receptor subunit mRNAs in the cellular processes of cultured cortical neurons and glia (69) 44

286

Qin, Z., see Yang, G.-Y. (69) 135 Qin, Z.-H., see Lammers, C.-H. (69) 281

Rance, N.E., see Abel, T.W. (69) 202

Sabel, B.A., see Kreutz, M.R. (69) 232 Sabel, M., see Sommer, C. (69) 53 Sautter, J., see Sommer, C. (69) 53 Schmidt, P., see Ciani, E. (69) 290 Schuller, J.J., Billings, L.M. and Marshall, J.F. Dopaminergic modulation of pallidal preproenkephalin mRNA (69) 149 Seidenbecher, C.I., see Kreutz, M.R. (69) 232 Sharma, M., see Pietruck, C. (69) 286 Shen, J., see Berezovska, O. (69) 273 Simerly, R.B., see Leng, N. (69) 73 Sommer, C., Sabel, M., Oertel, W.H., Kiessling, M. and Sautter, J. Temporo-spatial expression of bFGF and TGFβ2 in embryonic dopaminergic grafts in a rat model of Parkinson's disease (69)

Spindel, E.R., see Leng, N. (69) 73
Sumikawa, K., see Nishizaki, T. (69) 263
Sumikawa, K., see Nishizaki, T. (69) 295
Swartz, D., see Zhou, F.C. (69) 186
Sweerts, B.W., Jarrott, B. and Lawrence, A.J.
Expression of preprogalanin mRNA following acute and chronic restraint stress in brains of normotensive and hypertensive rats (69) 113

Spengler, D., see Ciani, E. (69) 290

Takeda, J., see Honma, Y. (69) 93
Tanaka, T., see Yukawa, K. (69) 124
Tanno, Y., see Honma, Y. (69) 93
Thi Man, N., see Wilkinson, F.L. (69) 10
Thomas, P., see Wilkinson, F.L. (69) 10
Tiao, N., see Feng, Z. (69) 144
Tischmeyer, W., see Kreutz, M.R. (69) 232
Tojo, M., see Honma, Y. (69) 93
Tonegawa, S., see Berezovska, O. (69) 273
Tsuji, S., see Yukawa, K. (69) 124
Tsukamoto, T., see Honma, Y. (69) 93

Ueda, H., see Inoue, M. (69) 302 Ulfhake, B., see Johnson, H. (69) 21

Vecchiola, A., Collyer, P., Figueroa, R.,
Labarca, R., Bustos, G. and Magendzo, K.
Differential regulation of μ-opioid receptor mRNA in the nucleus accumbens shell and core accompanying amphetamine behavioral sensitization (69) 1
Verity, A.N., see Baecker, P.A. (69) 209

Vorwerk, C.K., see Kreutz, M.R. (69) 232

Walton, M., see Woodgate, A. (69) 84
Wanaka, A., see Honma, Y. (69) 93
Wei, L.-N., see Hu, X. (69) 35
Wilkinson, F.L., Thi Man, N., Manilal, S.B.,
 Thomas, P., Neal, J.W., Harper, P.S., Jones,
 A.L. and Morris, G.E.
 Localization of rabbit huntingtin using a
 new panel of monoclonal antibodies (69) 10
Williams, F.G., see Ko, J.L. (69) 171
Woodgate, A., MacGibbon, G., Walton, M. and
 Dragunow, M.
 The toxicity of 6-hydroxydopamine on PC12
 and P19 cells (69) 84

Xie, G.-X., see Pietruck, C. (69) 286 Xu, Y., see Zhou, F.C. (69) 186

Yajima, S., see Lammers, C.-H. (69) 281
Yamada, K., see Nakaso, K. (69) 155
Yamada, T., see Inoue, M. (69) 302
Yanagawa, T., see Nakaso, K. (69) 155
Yang, G.-Y., Gong, C., Qin, Z., Liu, X.-H. and Lorris Betz, A.
Tumor necrosis factor alpha expression produces increased blood-brain barrier permeability following temporary focal cerebral ischemia in mice (69) 135
Yokoya, S., see Honma, Y. (69) 93

Yukawa, K., Tanaka, T., Tsuji, S. and Akira, S. Regulation of transcription factor C/ATF by the cAMP signal activation in hippocampal neurons, and molecular interaction of C/ATF with signal integrator CBP/p300 (69) 124

Zhou, F.C., Patel, T.D., Swartz, D., Xu, Y. and Kelley, M.R.
Production and characterization of an antiserotonin 1A receptor antibody which detects functional 5-HT<sub>1A</sub> binding sites (69) 186

Zhu, Q., see Levavasseur, F. (69) 104